

Artemisinin and Derivatives Cancer Studies

Cervical Cancer

- (1) Chen HH *et al.* Inhibition of human cancer cell line growth and human umbilical vein endothelial cell angiogenesis by artemisinin derivatives in vitro. *Pharmacol Res.* 2003 Sep; 48(3):231-6.
- (2) Disbrow GL *et al.* Dihydroartemisinin is cytotoxic to papillomavirus-expressing epithelial cells in vitro and in vivo. *Cancer Research.* 2005 Dec 1; 65(23):10854-61.
- (3) Ju HK *et al.* Standardized flavonoid-rich fraction of *Artemisia princeps* Pampanini cv. Sajabal induces apoptosis via mitochondrial pathway in human cervical cancer HeLa cells. *J Ethnopharmacol.* 2012 Mar 18
- (4) Paik IH *et al.* Second generation, orally active, antimalarial, artemisinin-derived trioxane dimers with high stability, efficacy, and anticancer activity. *J Med Chem.* 2006 May 4;49(9):2731-4.
- (5) Thanaketpaisarn O *et al.* Artesunate enhances TRAIL-induced apoptosis in human cervical carcinoma cells through inhibition of the NF- κ B and PI3K/Akt signaling pathways. *Int J Oncol.* 2011 Jul;39(1):279-85.
- (6) Zheng JS *et al.* Artesunate suppresses human endometrial carcinoma RL95-2 cell proliferation by inducing cell apoptosis [Article in Chinese]. *Nan Fang Yi Ke Da Xue Xue Bao.* 2008 Dec;28(12):2221-3.