

Artemisinin and Derivatives Cancer Studies

Colorectal Cancer

- (1) Cui C *et al.* Influence of artesunate and oxymatrine on immunosuppressive effect of mouse colon-rectal carcinoma cell line. *Xiandai Mianyixue* 26, 152-156, 2006.
- (2) Li LN *et al.* Differential sensitivity of colorectal cancer cell lines to artesunate is associated with expression of beta-catenin and E-cadherin. *Eur J Pharmacol.* 2008 Apr 3.
- (3) Li LN *et al.* Artesunate attenuates the growth of human colorectal carcinoma and inhibits hyperactive Wnt/beta-catenin pathway. *Int J Cancer.* 2007 Sep 15;121(6):1360-5.
- (4) Lu JJ *et al.* Characterization of dihydroartemisinin-resistant colon carcinoma HCT116/R cell line. *Mol Cell Biochem.* 2012 Jan;360(1-2):329-37.
- (5) Lu JJ *et al.* The anti-cancer activity of dihydroartemisinin is associated with induction of iron-dependent endoplasmic reticulum stress in colorectal carcinoma HCT116 cells. *Invest New Drugs.* 2010 Jul 7.
- (6) Riganti C *et al.* Artemisinin induces doxorubicin resistance in human colon cancer cells via calcium-dependent activation of HIF-1alpha and P-glycoprotein overexpression. *Br J Pharmacol.* 2009 Apr;156(7):1054-66.
- (7) Woerdenbag HJ *et al.* Cytotoxicity of flavonoids and sesquiterpene lactones from Arnica species against the GLC4 and the COLO 320 cell lines. *Planta Med* 60:434-437, 1994.
- (8) Zheng GQ. Cytotoxic terpenoids and flavonoids from *Artemisia annua*. *Planta Med* 60:54-57, 1994.